

New proofs of Rosenthal's ℓ^1 -theorem and the Josefson–Nissenzweig theorem

EHRHARD BEHREND

ABSTRACT. We give elementary proofs of the theorems mentioned in the title. Our methods rely on a simple version of Ramsey theory and a martingale difference lemma.

They also provide quantitative results: if a Banach space contains ℓ^1 only with a bad constant then every bounded sequence admits a subsequence which is “nearly” a weak Cauchy sequence.

1. INTRODUCTION

The aim of this paper is to give new, elementary proofs of the following theorems:

Rosenthal's ℓ^1 -theorem ([6]): Let (x_n) be a bounded sequence in a Banach space X . Either there is a subsequence which is equivalent to the ℓ^1 -basis or there is a subsequence (x_{n_k}) which is weakly Cauchy (i.e. $(x'(x_{n_k}))$ converges for every $x' \in X'$).

The Josefson–Nissenzweig theorem ([4], [5]): For every infinite-dimensional Banach space X there are $x'_1, x'_2, \dots$ in X' such that $\|x'_n\| = 1$ for all n , but $x'_n \rightarrow 0$ w.r.t. the weak*-topology.

They are treated in several textbooks (see e.g. [2] where the reader also may find some historical remarks). We are going to present new proofs which are more elementary than those in the literature. Also, in the case of Rosenthal's theorem, we are able to derive a quantitative variant which reads as follows for real spaces (cf. section 3 where also the complex case is discussed): If (x_n) is bounded, $\varepsilon > 0$, and if for every infinite $M \subset \mathbb{N}$ there are $i_1 < \dots < i_r$ in M and $a_1, \dots, a_r \in \mathbb{R}$ with $\sum |a_\rho| = 1$ and $\|\sum a_\rho x_{i_\rho}\| \leq \varepsilon$, then there is a subsequence (x_{n_k}) such that $\limsup_k x'(x_{n_k}) - \liminf_k x'(x_{n_k}) \leq 2\varepsilon$ for all x' with $\|x'\| = 1$.

We prepare the proof of this in *section 2* where we treat an easy-to-formulate Ramsey-type theorem (the proof of which is elementary but technical). In *section 4* we show that a slightly modified assumption in Rosenthal's theorem even gives rise to norm convergent subsequences; this has already been published in [1], the present argument, however, is much simpler.

Next, in *section 5*, we are going to prepare the proof of the Josefson–Nissenzweig theorem (the section contains some facts about Banach limits and a simple martingale lemma). *Section 6* uses the circle of ideas presented in *section 3* to prove the Josefson–Nissenzweig theorem. This idea is also fundamental in [2], our proof however is more direct and gives a slightly sharper result.

2. AN ELEMENTARY RAMSEY THEOREM

Theorem 2.1 *For $r \in \mathbb{N}$ let T_r be a family of r -tuples of increasing integers. Suppose that*

$$(1) \quad \forall_{\substack{M \subset \mathbb{N} \\ M \text{ infinite}}} \exists_{\substack{i_1, i_2, \dots \in M \\ i_1 < i_2 < \dots}} \forall_r (i_1, \dots, i_r) \in T_r.$$

Then it follows that

$$(2) \quad \exists_{\substack{M_0 \subset \mathbb{N} \\ M_0 \text{ infinite}}} \forall_r \exists_{\substack{i_1, i_2, \dots \in M_0 \\ i_1 < \dots < i_r}} (i_1, \dots, i_r) \in T_r.$$

The rest of this section is devoted to the proof of 2.1. The key will be the following

Definition 2.2 Let $M \subset \mathbb{N}$ be infinite, $i_1, \dots, i_k \in \mathbb{N}$, $i_1 < \dots < i_k$. Moreover let $(T_r)_{r=1,2,\dots}$ be as in 2.1.

- (i) $i_1, \dots, i_k \downarrow M$ abbreviates the following fact:
If $i_{k+1} < i_{k+2} < \dots$ are points in M with $i_{k+1} > i_k$, then there is an r such that $(i_1, \dots, i_r) \notin T_r$.
The case $k = 0$ will also be admissible, we will then write $\emptyset \downarrow M$.
- (ii) $i_1, \dots, i_k \uparrow M$ stands for the following:
Whenever N is an infinite subset of M , there are $i_{k+1} < i_{k+2} < \dots$ in N with $i_{k+1} > i_k$ such that $(i_1, \dots, i_r) \in T_r$ for every r .
Again the definition is meant to contain the case $k = 0$.

Note. Our “ $\downarrow$ ” and “ $\uparrow$ ” are closely related with “acceptance” and “rejection” in Diestel’s book ([2], p. 192). Our approach, however, is more direct since we only have in mind a special version of a Ramsey theorem.

The following facts are immediate consequences of the definitions, they are stated only for the sake of easy reference.

Observation 2.3

- (i) (1) of the theorem just means $\emptyset \uparrow \mathbb{N}$, and (2) follows as soon as one has found an M_0 such that $i_1, \dots, i_k \uparrow M_0$ for arbitrary $i_1 < \dots < i_k$ in M_0 .
- (ii) Let $i_1 < \dots < i_k$ be given and $M \subset \mathbb{N}$ be infinite. If $i_1, \dots, i_k \downarrow M$ does not hold, then there are $i_{k+1} < i_{k+2} < \dots$ (with $i_{k+1} > i_k$) in M such that $(i_1, \dots, i_r) \in T_r$ for every r .
- (iii) If $i_1, \dots, i_k \downarrow M$ and $N \subset M$ is infinite, then $i_1, \dots, i_k \downarrow N$. The same holds if the “ $\downarrow$ ” are replaced by “ $\uparrow$ ”.

Here is the first step of our construction:

Lemma 2.4 *There is an infinite $\widetilde{M}_0 \subset \mathbb{N}$ such that*

$$(3) \quad i_1, \dots, i_k \downarrow \widetilde{M}_0 \quad \text{or} \quad i_1, \dots, i_k \uparrow \widetilde{M}_0$$

for each choice of $i_1 < \dots < i_k$ in $\widetilde{M}_0$ (including the case $k = 0$).

Proof. It will be convenient to say that an infinite $\widetilde{M}_0$ satisfies $(3)_s$ (where $s \in \mathbb{N}_0$) if the assertion (3) holds under the additional assumption that $\{i_1, \dots, i_k\}$ is contained in the set of the first s elements of $\widetilde{M}_0$. We combine the following observations:

- We are looking for an infinite $\widetilde{M}_0$ such that $\widetilde{M}_0$ satisfies $(3)_s$ for every s .
- Suppose we are able to make the following induction work: Given an infinite $\widetilde{M}^{(s)}$ (which we write in increasing order as $\widetilde{M}^{(s)} = \{i_1, i_2, \dots, i_s, \dots\}$) such that $(3)_s$ holds for $\widetilde{M}^{(s)}$, there is an infinite subset N of $\{i_{s+1}, i_{s+2}, \dots\}$ such that $\widetilde{M}^{(s+1)} := \{i_1, \dots, i_s\} \cup N$ satisfies $(3)_{s+1}$.

This would suffice: We start our construction by setting $\widetilde{M}^{(0)} := \mathbb{N}$ (note that $\widetilde{M}^{(0)}$ satisfies $(3)_0$ by 2.3(i)), use the induction to

construct the $\widetilde{M}^{(0)} \supset \widetilde{M}^{(1)} \supset \widetilde{M}^{(2)} \dots$ and set $\widetilde{M}_0 =$ “the collection of the s ’th elements of $\widetilde{M}^{(s)}$, $s \in \mathbb{N}$.” For fixed s , $\widetilde{M}_0$ has the same first s elements as $\widetilde{M}^{(s)}$, so that in view of 2.3(iii) $(3)_s$ necessarily holds for $\widetilde{M}_0$.

Therefore let’s concentrate on the induction step. Let $s \geq 0$ and $\widetilde{M}^{(s)}$ with $(3)_s$ be given. Denote by $\Delta_1, \dots, \Delta_{2^s}$ the 2^s different subsets of $\{i_1, \dots, i_s\}$.

We will construct infinite subsets $\widetilde{N}^{[1]} \supset \widetilde{N}^{[2]} \supset \dots \supset \widetilde{N}^{[2^s]}$ of $\{i_{s+2}, \dots\}$ such that either $\Delta_j, i_{s+1} \uparrow \widetilde{N}^{[j]}$ or $\Delta_j, i_{s+1} \downarrow \widetilde{N}^{[j]}$ for every j . In view of 2.3(iii) it is then clear that $\widetilde{M}^{(s+1)} := \{i_1, \dots, i_{s+1}\} \cup \widetilde{N}^{[2^s]}$ has $(3)_{s+1}$.

First consider Δ_1, i_{s+1} . Either we have $\Delta_1, i_{s+1} \downarrow \widetilde{M}^{(s)}$ (in which case we put $\widetilde{N}^{[1]} := \{i_{s+2}, \dots\}$) or there is an infinite subset $\widetilde{N}^{[1]}$ of $\{i_{s+2}, \dots\}$ such that $\Delta_1, i_{s+1} \uparrow \widetilde{N}^{[1]}$ (see 2.3(ii)).

Secondly, we investigate Δ_2, i_{s+1} . Either $\Delta_2, i_{s+1} \downarrow \widetilde{N}^{[1]}$ (we will put $\widetilde{N}^{[2]} := \widetilde{N}^{[1]}$ in this case) or there is an infinite subset $\widetilde{N}^{[2]}$ of $\widetilde{N}^{[1]}$ such that $\Delta_2, i_{s+1} \uparrow \widetilde{N}^{[2]}$. It should be clear how to construct the remaining $\widetilde{N}^{[3]} \supset \dots \supset \widetilde{N}^{[2^s]}$.

In order to get an M_0 with (2) from $\widetilde{M}_0$ we need

Lemma 2.5 *Let $i_1 < \dots < i_k$ in $\widetilde{M}_0$ be given and suppose that $i_1, \dots, i_k \uparrow \widetilde{M}_0$. Then there are only finitely many $i > i_k$ in $\widetilde{M}_0$ such that $i_1, \dots, i_k, i \downarrow \widetilde{M}_0$.*

Proof. Suppose that this were not the case. Put $N =$ the collection of these i . N is infinite, and by $i_1, \dots, i_k \uparrow \widetilde{M}_0$ there would be $i_{k+1} < i_{k+2} < \dots$ in N (with $i_{k+1} > i_k$) such that $(i_1, \dots, i_r) \in T_r$ for every r . Note that this would contradict $i_1, \dots, i_{k+1} \downarrow \widetilde{M}_0$.

Finally, we are ready for the

Proof of theorem 2.1. We have already noted that (2) just means $i_1, \dots, i_k \uparrow M_0$ for $i_1 < \dots < i_k$ in M_0 . Similarly to the proof of 2.4 we introduce $(2)_s$ for $s \geq 0$: This is (2) with the same additional assumption as in $(3)_s$.

The construction parallels that of 2.4: We need an M_0 with $(2)_s$ for all s , we know that $M^{(0)} := \mathbb{N}$ satisfies $(2)_0$, and, given an $M^{(s)}$ with $(2)_s$, we only have to construct an $M^{(s+1)}$ with $(2)_{s+1}$. In this construction the first s elements of $M^{(s)}$ and $M^{(s+1)}$ should be identical; $M_0 =$ “the collection of the s ’th elements of $M^{(s)}$, $s \in \mathbb{N}$ ” then will have the desired properties.

Here is the induction. Write $M^{(s)} = \{i_1, \dots, i_s, i_{s+1}, \dots\}$ and put $N := \{i_{s+1}, \dots\}$. Since $\Delta \uparrow N$ for every $\Delta \subset \{i_1, \dots, i_s\}$ by assumption we conclude from 2.5 that there are only finitely many i in N such that $\Delta, i \downarrow N$

for any Δ . Choose $\tilde{N} \subset N$ such that $\tilde{N}$ does not contain such i . Then $M^{(s+1)} := \{i_1, \dots, i_s\} \cup \tilde{N}$ satisfies (2)_{s+1}, and this completes the proof.

3. A QUANTITATIVE VERSION OF ROSENTHAL'S ℓ^1 -THEOREM

To begin with, we restate a definition from [1].

Definition 3.1 Let (x_n) be a bounded sequence in a Banach space X , and $\varepsilon > 0$. We say that (x_n) admits ε - ℓ^1 -blocks if for every infinite $M \subset \mathbb{N}$ there are $a_1, \dots, a_r \in \mathbb{K}$ with $\sum |a_r| = 1$ and $i_1 < \dots < i_r$ in M such that $\|\sum a_\rho x_{i_\rho}\| \leq \varepsilon$.

Clearly there will be no subsequence of (x_n) equivalent to the ℓ^1 -basis iff (x_n) admits ε - ℓ^1 -blocks for arbitrarily small $\varepsilon > 0$. Thus Rosenthal's theorem is the assertion that (x_n) has a weak Cauchy subsequence provided it admits ε - ℓ^1 -blocks for all ε . Here is our quantitative version of this fact in the case of real spaces:

Theorem 3.2 *Let X be a real Banach space and (x_n) a bounded sequence. Suppose that, for some $\varepsilon > 0$, (x_n) admits small ε - ℓ^1 -blocks. Then there is a subsequence (x_{n_k}) of (x_n) such that (x_{n_k}) is “close to being a weak Cauchy sequence” in the following sense:*

$$\limsup x'(x_{n_k}) - \liminf x'(x_{n_k}) \leq 2\varepsilon$$

for every x' with $\|x'\| = 1$.

Remark. It is simple to derive the original theorem from 3.2. (If (x_n) and thus every subsequence has ε - ℓ^1 -blocks for all ε , apply 3.2 successively with ε running through a sequence tending to zero. The diagonal sequence which is obtained from this construction will be a weak Cauchy sequence.)

Proof. Suppose the theorem were not true. We claim that without loss of generality we may assume that there is a $\delta > 0$ such that

$$(4) \quad \varphi((x_{n_k})) := \sup_{\|x'\|=1} \left(\limsup_k x'(x_{n_k}) - \liminf_k x'(x_{n_k}) \right) > 2\varepsilon + \delta$$

for all subsequences (x_{n_k}) . In fact, if every subsequence contained another subsequence with a φ -value arbitrarily close to 2ε , an argument as in the

preceding remark would even provide one where $\varphi((x_{n_k})) \leq 2\varepsilon$ in contrast to our assumption.

Fix a $\tau > 0$ which will be specified later. The essential tool in order to get a contradiction will be the

Lemma *After passing to a subsequence we may assume that (x_n) satisfies the following conditions:*

- (i) *If C and D are finite disjoint subsets of $\mathbb{N}$ there are a $\lambda_0 \in \mathbb{R}$ and an $x' \in X'$ with $\|x'\| = 1$ such that $x'(x_n) < \lambda_0$ for $n \in C$ and $x'(x_n) > \lambda_0 + 2\varepsilon + \delta$ for $n \in D$.*
- (ii) *There are $i_1 < \dots < i_r$ in $\mathbb{N}$, $a_1, \dots, a_r \in \mathbb{R}$ with*

$$\sum |a_\rho| = 1, \quad \left| \sum a_\rho \right| \leq \tau, \quad \left\| \sum a_\rho x_{i_\rho} \right\| \leq \varepsilon.$$

Proof of the lemma. (i) Define, for $r \in \mathbb{N}$, T_r to be the collection of all $(i_1, \dots, i_r)$ (with $i_1 < \dots < i_r$) such that there are a $\lambda_0 \in \mathbb{R}$ and a normalized x' such that $x'(x_{i_\rho}) < \lambda_0$ if ρ is even and $> \lambda_0 + 2\varepsilon + \delta$ otherwise. (4) implies that (1) of 2.1 is valid. Thus there is an M_0 for which all $(i_1, \dots, i_r)$ are in T_r for $i_1 < \dots < i_r$ in M_0 . Let us assume that $M_0 = \mathbb{N}$.

Let C and D be finite disjoint subsets of $2\mathbb{N} = \{2, 4, \dots\}$. We may select $i_1 < \dots < i_r$ in $\mathbb{N}$ such that $C \subset \{i_\rho \mid \rho \text{ even}\}$ and $D \subset \{i_\rho \mid \rho \text{ odd}\}$. Because of $(i_1, \dots, i_r) \in T_r$ we have settled (i) provided C and D are in $2\mathbb{N}$, and all what's left to do is to consider (x_{2n}) instead of (x_n) .

(ii) By assumption we find $i_1 < \dots < i_r$, $a_1, \dots, a_r \in \mathbb{R}$ such that $\sum |a_\rho| = 1$ and $\left\| \sum a_\rho x_{i_\rho} \right\| \leq \varepsilon$ with arbitrarily large i_1 . Therefore we obtain $i_1^1 < \dots < i_{r_1}^1 < i_1^2 < \dots < i_{r_2}^2 < i_1^3 < \dots < i_{r_3}^3 < \dots$ and associated a_ρ^j . The numbers $\eta_j := \sum_{\rho=1}^{r_j} a_\rho^j$ all lie in $[-1, +1]$ so that we find $j < k$ with $|\eta_j - \eta_k| \leq 2\tau$. Let $i_1 < \dots < i_r$ be the family $i_1^j < \dots < i_{r_j}^j < i_1^k < \dots < i_{r_k}^k$, and define the $a_1, \dots, a_r$ by $\frac{1}{2}a_1^j, \dots, \frac{1}{2}a_{r_j}^j, -\frac{1}{2}a_1^k, \dots, -\frac{1}{2}a_{r_k}^k$.

We are now ready to derive a contradiction. On the one hand, by (ii) of the lemma, we find $i_1 < \dots < i_r$ in $\mathbb{N}$, $a_1, \dots, a_r \in \mathbb{R}$, $\sum |a_\rho| = 1$, $\left| \sum a_\rho \right| \leq \tau$ with $\left\| \sum a_\rho x_{i_\rho} \right\| \leq \varepsilon$. On the other hand we may apply (i) with $C := \{i_\rho \mid a_\rho < 0\}$, $D := \{i_\rho \mid a_\rho > 0\}$. We put $\alpha := -\sum_{\rho \in C} a_\rho$, $\beta := \sum_{\rho \in D} a_\rho$, and we note that $|\alpha - \beta| \leq \tau$, $\alpha + \beta = 1$ so that $|\beta - \frac{1}{2}| \leq \tau$; hence

$$\varepsilon \geq \left\| \sum a_\rho x_{i_\rho} \right\| \geq \sum a_\rho x'(x_{i_\rho}) \geq -\lambda_0 \alpha + (\lambda_0 + 2\varepsilon + \delta) \beta \geq -|\lambda_0| \tau + \varepsilon + \frac{\delta}{2} - \tau \delta.$$

This expression can be made larger than ε if τ has been chosen sufficiently small (note that the numbers $|\lambda_0|$ are bounded by $\sup \|x_n\|$), a contradiction which proves the theorem.

Note. Since for the unit vector basis (x_n) of real ℓ^1 the assumption of the theorem holds with $\varepsilon = 1$ and since for every subsequence (x_{n_k}) one may find $\|x'\| = 1$ with $\limsup x'(x_{n_k}) - \liminf x'(x_{n_k}) = 2$ there can be no better constant than that given in our theorem.

Let's now turn to the *complex case* which has to be treated in a slightly different way. The quantitative version of Rosenthal's ℓ^1 -theorem here reads as follows:

Theorem 3.3 *Let (x_n) be a sequence in a complex Banach space X such that, for some $\varepsilon > 0$, (x_n) admits ε - ℓ^1 -blocks. Then there is a subsequence (x_{n_k}) such that, for x' with $\|x'\| = 1$, the diameter of the set of accumulation points of $(x'(x_{n_k}))_k$ is at most $8\varepsilon/\sqrt{2}$.*

Proof. The proof is similar to the preceding one. Again we know that – if the theorem does not hold – the numbers

$$\sup_{\|x'\|=1} \{\text{diameter of the accumulation points of } (x'(x_{n_k}))\}$$

are greater than $8\varepsilon/\sqrt{2} + \delta$ for a suitable $\delta > 0$ and all (x_{n_k}) , and again we fix a $\tau > 0$.

The key is this time the

Lemma *Without loss of generality the sequence (x_n) satisfies the following conditions.*

- (i) *Whenever C and D are disjoint and finite subsets of $\mathbb{N}$ there are $z_0, w_0 \in \mathbb{C}$ with $|w_0| \geq 8\varepsilon/\sqrt{2} + \delta$ and an x' with $\|x'\| = 1$ such that $|x'(x_n) - z_0| \leq \tau$ for $n \in C$ and $|x'(x_n) - (z_0 + w_0)| \leq \tau$ for $n \in D$.*
- (ii) *We may assume that the a 's of 3.1 additionally satisfy $|\sum a_\rho| \leq \tau$.*

The proof is similar and is therefore omitted.

Now let $S_1, \dots, S_4$ be the quadrants in the complex plane, i.e., S_j is the collection of those numbers whose arguments lie between $(j-1)\pi/2$ and $j\pi/2$. We need the following obvious facts:

- If $\sum |a_\rho| = 1$ there is a j such that $\sum_{a_\rho \in S_j} |a_\rho| \geq 1/4$.

- If $\sum_{a_\rho \in S_j} |a_\rho| \geq 1/4$, then $\sum_{a_\rho \in S_j} |a_\rho| \geq \sqrt{2}/8$.

Now choose $i_1 < \dots < i_r$, $a_1, \dots, a_r$ with $\sum |a_r| = 1$, $|\sum a_\rho| \leq \tau$, $|\sum a_\rho x_{i_\rho}| \leq \varepsilon$. Write $\{i_1, \dots, i_r\}$ as the disjoint union of C and D , where D contains the i_ρ with $a_\rho \in S_1$ (without loss of generality we assume that $\sum_{a_\rho \in S_1} |a_\rho| \geq 1/4$).

With z_0, w_0, x' as in (i) of the lemma it follows that

$$\begin{aligned}
\varepsilon &\geq \left\| \sum a_\rho x_{i_\rho} \right\| \geq \left| \sum a_\rho x'(x_{i_\rho}) \right| \\
&\geq \left| \sum_{a_\rho \in S_1} a_\rho (z_0 + w_0) + \sum_{a_\rho \notin S_1} a_\rho z_0 \right| - \tau \sum |a_\rho| \\
&= \left| \sum a_\rho z_0 + \sum_{a_\rho \in S_1} a_\rho w_0 \right| - \tau \\
&\geq |w_0| \left| \sum_{a_\rho \in S_1} a_\rho \right| - \tau(1 + |z_0|) \\
&\geq |w_0| \frac{\sqrt{2}}{8} - \tau(1 + |z_0|) \\
&\geq \varepsilon + \frac{\sqrt{2}}{8} \delta - \tau(1 + |z_0|),
\end{aligned}$$

and this yields a contradiction for small τ .

Note. It seems to be difficult to decide whether $8\varepsilon/\sqrt{2}$ can in general be replaced by a better estimate.

4. BLOCKINGS WITH UNIFORMLY BOUNDED LENGTH

The following result has been published in [1]. Here we present a much simpler proof which uses theorem 2.1.

Theorem 4.1 *Let (x_n) be a bounded sequence in a real or complex Banach space X such that there is an $r \in \mathbb{N}$ with the following property: Whenever $M \subset \mathbb{N}$ is infinite and $\varepsilon > 0$, there are $i_1 < \dots < i_r$ in M and $a_1, \dots, a_r \in \mathbb{K}$ with $\sum |a_\rho| = 1$ such that $\|\sum a_i x_{i_\rho}\| \leq \varepsilon$. Then (x_n) has a convergent subsequence.*

Proof. Fix $\varepsilon > 0$. The first step is as in [1], we refer the reader to this paper: One can choose the same a 's for all M . Fix $a_1, \dots, a_r$ and suppose that $a_r \neq$

0. Define sets of $\tilde{r}$ -tupels $T_{\tilde{r}}$ as follows. $T_{\tilde{r}}$ is the collection of *all* $\tilde{r}$ -tupels if $\tilde{r} \neq r$, and the set of those $(i_1, \dots, i_r)$ with $i_1 < \dots < i_r$ and $\|\sum a_\rho x_{i_\rho}\| \leq \varepsilon$ if $\tilde{r} = r$. Then 2.1 may be applied, and we get (x_{n_k}) such that $\|\sum a_\rho x_{n_{k_\rho}}\| \leq \varepsilon$ for arbitrary $k_1 < \dots < k_r$. In particular $x_{n_r}, x_{n_{r+1}}, x_{n_{r+2}}, \dots$ lie in the ball with center $-\sum_1^{r-1} (a_\rho/a_r) x_{n_\rho}$ and radius $\varepsilon/|a_r|$. Starting this construction with $\varepsilon = 1$ and applying it repeatedly to $\varepsilon = 1/2, \varepsilon = 1/3, \dots$ provides a descending family of subsequences for which the diagonal sequence surely is convergent.

Note. In this situation a quantitative version is not to be expected in general. Consider e.g. the unit vector basis (e_n) in c_0 . For $\varepsilon > 0$ fixed one can find the $a_1, \dots, a_r$ with the same r for all M , but $\|e_n - e_m\| = 2$ for $n \neq m$.

5. THE JOSEFSON–NISSENZWEIG THEOREM: PREPARATIONS

Lemma 5.1 *For $\ell = 1, 2, \dots$ and $\varepsilon_1, \dots, \varepsilon_\ell \in \{0, 1\}$ let $r_{\varepsilon_1 \dots \varepsilon_\ell}$ be a number such that the family $((r_{\varepsilon_1 \dots \varepsilon_\ell})_{\varepsilon_1 \dots \varepsilon_\ell})_{\ell=1,2,\dots}$ satisfies $|r_{\varepsilon_1 \dots \varepsilon_\ell}| \leq 2^{-\ell}$ and $r_{\varepsilon_1 \dots \varepsilon_\ell} = r_{\varepsilon_1 \dots \varepsilon_\ell 0} + r_{\varepsilon_1 \dots \varepsilon_\ell 1}$ for all $\ell, \varepsilon_1 \dots \varepsilon_\ell$. Define*

$$\eta_\ell := \sum_{\varepsilon_1 \dots \varepsilon_\ell} (r_{\varepsilon_1 \dots \varepsilon_\ell 0} - r_{\varepsilon_1 \dots \varepsilon_\ell 1}).$$

Then $\sum |\eta_\ell|^2 \leq 1$ so that in particular $\eta_\ell \rightarrow 0$.

Proof. An elegant proof could be given using martingales: the $r_{\varepsilon_1 \dots \varepsilon_\ell}$ give rise to a bounded martingale, the martingale convergence theorem guarantees the existence of a limit f in L^1 , and the η_ℓ are the integrals over f multiplied by suitable Rademacher functions.

However, a much simpler approach is possible. Set

$$\begin{aligned} a_\ell &:= 2^\ell \sum_{\varepsilon_1 \dots \varepsilon_\ell} |r_{\varepsilon_1 \dots \varepsilon_\ell}|^2 \\ b_\ell &:= 2^\ell \sum_{\varepsilon_1 \dots \varepsilon_\ell} |r_{\varepsilon_1 \dots \varepsilon_\ell 0} - r_{\varepsilon_1 \dots \varepsilon_\ell 1}|^2 \\ c_\ell &:= \sum_{\varepsilon_1 \dots \varepsilon_\ell} |r_{\varepsilon_1 \dots \varepsilon_\ell 0} - r_{\varepsilon_1 \dots \varepsilon_\ell 1}|. \end{aligned}$$

Then $a_{\ell+1} - a_\ell = b_\ell$ (by the parallelogram law $|\alpha + \beta|^2 + |\alpha - \beta|^2 = 2(|\alpha|^2 + |\beta|^2)$) so that the a_ℓ are increasing. Surely $a_\ell \leq 1$, and we get $\sum b_\ell \leq 1$.

Finally note that $|\eta_\ell| \leq c_\ell$ and that $c_\ell^2 \leq b_\ell$ since

$$(|\alpha_1| + \cdots + |\alpha_k|)^2 \leq k(|\alpha_1|^2 + \cdots + |\alpha_k|^2) \quad \text{for all families } \alpha_1, \dots, \alpha_k.$$

Next we have to remind the reader of the definition of a Banach limit which can be found in nearly every textbook on functional analysis. A *Banach limit* is an $L \in (\ell^\infty)'$ such that $L((1, 1, \dots)) = 1 = \|L\|$, and $L((x_1, x_2, \dots)) = L((x_2, x_3, \dots))$ for every (x_n) . It is known that such L exist. Some elementary properties will be important for us:

Lemma 5.2 *Define $\lambda_\ell, \mu_{\varepsilon_1 \dots \varepsilon_\ell} \in \ell^\infty$ for $\ell = 1, 2, \dots$ and $\varepsilon_1, \dots, \varepsilon_\ell \in \{0, 1\}$ as follows:*

$$\begin{aligned} \lambda_1 &= (1, -1, 1, -1, \dots) \\ \lambda_2 &= (1, 1, -1, -1, 1, 1, \dots) \\ \lambda_3 &= (1, 1, 1, 1, -1, -1, -1, -1, \dots) \\ &\vdots \\ \mu_0 &= (1, 0, 1, 0, \dots), \quad \mu_1 = (0, 1, 0, 1, \dots) \\ \mu_{00} &= (1, 0, 0, 0, 1, 0, 0, 0, 1, \dots), \quad \mu_{01} = (0, 0, 1, 0, 0, 0, 1, 0, \dots) \\ \mu_{10} &= (0, 1, 0, 0, 0, 1, 0, 0, 0, \dots), \quad \mu_{11} = (0, 0, 0, 1, 0, 0, 0, 1, \dots); \end{aligned}$$

in general: $\mu_{\varepsilon_1 \dots \varepsilon_\ell}$ is 1 at positions of the form $k \cdot 2^\ell + 1 + \varepsilon_1 2^0 + \cdots + \varepsilon_\ell 2^{\ell-1}$ ($k = 0, 1, \dots$) and 0 otherwise, and

$$\lambda_\ell = \sum_{\varepsilon_1, \dots, \varepsilon_{\ell-1}} (\mu_{\varepsilon_1, \dots, \varepsilon_{\ell-1}, 0} - \mu_{\varepsilon_1, \dots, \varepsilon_{\ell-1}, 1}).$$

Then $|L(\mu_{\varepsilon_1 \dots \varepsilon_\ell} x)| \leq 2^{-\ell}$ for every $x \in \ell^\infty$ with $\|x\| \leq 1$.

Proof. Let $T : \ell^\infty \rightarrow \ell^\infty$ be the shift operator $(y_1, y_2, \dots) \mapsto (0, y_1, y_2, \dots)$ and x_0 the pointwise product of $\mu_{\varepsilon_1 \dots \varepsilon_\ell}$ with x . Then $L(T\tilde{x}) = L(\tilde{x})$ for every $\tilde{x}$, and $\|x_0 + Tx_0 + T^2x_0 + \cdots + T^{2^\ell-1}x_0\| \leq 1$. Hence $2^\ell |L(x_0)| \leq 1$.

6. THE JOSEFSON–NISSENZWEIG THEOREM

Theorem 6.1 ([4], [5]) *A Banach space X is either finite-dimensional or there exist normalized $x'_1, x'_2, \dots$ such that $x'_n(x) \rightarrow 0$ for every x .*

Proof.

CASE 1: ℓ^1 is not contained in X' . Suppose that every weak*-convergent sequence is already norm convergent; we will show that X' is finite-dimensional.

Let (x'_n) be a bounded sequence. By Rosenthal's theorem and since ℓ^1 does not embed into X' we find a subsequence which is weakly Cauchy and thus weak*-convergent. By our assumption it is convergent, and thus X' is finite-dimensional.

CASE 2: ℓ^1 embeds into X' , i.e., there are x'_n in X' and $A, B > 0$ such that

$$A \sum_1^r |t_i| \geq \left\| \sum_1^r t_i x'_i \right\| \geq B \sum_1^r |t_i|$$

for arbitrary r and $t_1, \dots, t_r \in \mathbb{K}$.

In order to continue we remind the reader of the following notion: A sequence (y'_n) is said to be obtained from the (x'_n) by *blocking* if there are disjoint finite sets $A_1, A_2, \dots$ in $\mathbb{N}$ with $A_1 \leq A_2 \leq \dots$ and numbers (a_k) with $\sum_{k \in A_n} |a_k| = 1$ for every n such that $y'_n = \sum_{k \in A_n} a_k x'_k$. Note in particular that all subsequences arise in this way.

CASE 2.1. It is possible to get (y'_n) by blocking (x_n) such that $y'_n \rightarrow 0$ with respect to the weak*-topology. Then we are done since $\|y'_n\| \geq B$ so that the $y'_n/\|y'_n\|$ have the desired properties.

CASE 2.2. For no blocking (y'_n) we have $y'_n \rightarrow 0$ (w.r.t. the weak*-topology). In order to measure the property of being a weak*-null sequence we introduce the number

$$\varphi((y'_n)) := \sup_{\|x\|=1} \limsup |y'_n(x)|$$

for the (y'_n) constructed as before. In the case under consideration we know that always $\varphi((y'_n)) > 0$. We claim that even more is true.

CLAIM 1. There are a $\delta > 0$ and a block sequence (y'_n) such that $\varphi((z'_n)) = \delta$ for every (z'_n) which is obtained from (y'_n) by blocking.

Proof of claim 1. Let $\delta_0 \geq 0$ be the infimum of the numbers $\varphi((y'_n))$, where the infimum runs over all block sequences (y'_n) . Choose $(y_n^{[1]})$, a block sequence of (x'_n) , such that $\varphi((y_n^{[1]})) \leq \delta_0 + 1/2^0$. Let δ_1 be the infimum of the $\varphi((y'_n))$, where this time only blockings of $(y_n^{[1]})$ are under consideration. Since a block sequence of a block sequence is a block sequence we have

$\delta_0 \leq \delta_1$. Choose $(y_n^{[2]})$, a block sequence of $(y_n^{[1]})$, with $\varphi((y_n^{[2]})) \leq \delta_1 + 1/2^1$. In this way we get successively $(y_n^{[1]})$, $(y_n^{[2]})$, $\dots$, and $\delta_1 \leq \delta_2 \leq \dots$, where $(y_n^{[k+1]})$ is obtained from $(y_n^{[k]})$ by blocking and where $\delta_k \leq \varphi((y_n^{[k]})) \leq \delta_k + 1/2^k$ for all block sequences (y_n') of $(y_n^{[k+1]})$. Our candidate is the diagonal sequence (y_n') containing the n 'th element of the $(y_n^{[n]})$ for every n . For every k , (y_n') is – after possibly finitely many exceptions – a block sequence of $(y_n^{[k]})$; therefore the φ -value lies between δ_{k+1} and $\delta_k + 1/2^k$. It follows that $\delta := \sup \delta_k$ has the claimed properties; note that we also know that $\delta > 0$ since $\delta = \varphi((y_n'))$.

CLAIM 2. Fix (y_n') and $\delta > 0$ as in claim 1. Further let $\tau > 0$ be arbitrary. There is a subsequence which we will denote by (z_n') with the following property: It is possible to find normalized $x_1, x_2, \dots$ in X such that:

$$\begin{aligned} |z_n'(x_1) - \delta| &\leq \tau \text{ for all } n; \\ |z_n'(x_2) - \delta| &\leq \tau \text{ for } n = 3, 5, \dots \\ |z_n'(x_2) + \delta| &\leq \tau \text{ for } n = 4, 6, \dots \\ |z_n'(x_3) - \delta| &\leq \tau \text{ for } n = 5, 6, 9, 10, \dots \\ |z_n'(x_3) + \delta| &\leq \tau \text{ for } n = 7, 8, 11, 12, \dots \\ &\vdots \end{aligned}$$

(In general $z_n'(x_k)$ is τ -close to δ on segments of length 2^{k-2} beginning at $2^{k-1} + 1, 2 \cdot 2^{k-1} + 1, 3 \cdot 2^{k-1} + 1, \dots$ and τ -close to $-\delta$ at segments of the same length, beginning at $2^{k-1} + 2^{k-2} + 1, 2 \cdot 2^{k-1} + 2^{k-2} + 1, 3 \cdot 2^{k-1} + 2^{k-2} + 1$; thus, if we regard the x_k as functions on the set $\{z_n' \mid n \in \mathbb{N}\}$ they behave like the Rademacher functions, at least for large n .)

Proof of claim 2. By assumption we know that $\varphi((y_n')) = \delta$, and this makes it easy to find x_1 with $\|x_1\| = 1$ and a subsequence $(w_n^{[1]})$ of (y_n') with $|w_n^{[1]}(x_1) - \delta| \leq \tau$ for every n . Our final (z_n') will be a subsequence of $(w_n^{[1]})$ so that we will have no problems with x_1 . Put $z_1' := w_1^{[1]}$, $z_2' := w_2^{[2]}$. Now consider

$$(u_n) := \left(\frac{w_3^{[1]} - w_4^{[1]}}{2}, \frac{w_5^{[1]} - w_6^{[1]}}{2}, \dots \right).$$

This is a block sequence of (y_n') so that $\varphi((u_n)) = \delta$. Hence we find a normalized x_2 such that for infinitely many n , say $n \in N$, we have $|u_n(x_2) - \delta| \leq \tau'$; here τ' denotes any positive number such that

$$|\alpha|, |\beta| \leq \delta + \tau', \quad \left| \frac{1}{2}(\alpha + \beta) - \delta \right| \leq \tau' \quad \Rightarrow \quad |\alpha - \delta| \leq \tau, \quad |\beta - \delta| \leq \tau.$$

Since $\limsup |w_n^{[1]}(x_2)| \leq \delta$ we may also assume that for $n \in N$ and $u_n = (w_{2n-1}^{[1]} - w_{2n}^{[1]})/2$, the w 's satisfy $|w_{2n-1}^{[1]}(x_2)|, |w_{2n}^{[1]}(x_2)| \leq \delta + \tau'$. Thus $|w_{2n-1}^{[1]}(x_2) - \delta| \leq \tau$ and $|w_{2n}^{[1]}(x_2) + \delta| \leq \tau$.

Let $(w_n^{[2]})$ be the sequence of the $w_{2n-1}^{[1]}, w_{2n}^{[1]}$ with $n \in N$. Set $z'_3 := w_2^{[2]}$, $z'_4 := w_3^{[2]}$.

We consider now

$$\left(\frac{w_1^{[2]} + w_2^{[2]} - w_3^{[2]} - w_4^{[2]}}{4}, \frac{w_5^{[2]} + w_6^{[2]} - w_7^{[2]} - w_8^{[2]}}{4}, \dots \right) =: (u_n).$$

Again we find an infinite N and an x_3 such that $|u_n(x_3) - \delta| \leq \tau'$, where this time $\tau' > 0$ is such that

$$|\alpha_1|, |\alpha_2|, |\alpha_3|, |\alpha_4| \leq \delta + \tau' \text{ and } \left| \frac{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4}{4} - \delta \right| \leq \tau'$$

$$\Rightarrow |\alpha_j - \delta| \leq \tau \text{ for } j = 1, 2, 3, 4.$$

Also we can assume that the $|w_{4n-j}^{[2]}(x_3)| \leq \delta + \tau'$ for $n \in N$, $j = 0, 1, 2, 3$. Let $(w^{[3]})$ consist of the $w_{4n-3}^{[2]}, \dots, w_{4n}^{[2]}$ with $n \in N$, and define $z'_5, \dots, z'_8$ to be $w_1^{[3]}, \dots, w_4^{[3]}$ respectively.

It should be clear how this construction (which is similar to that in [3]) has to be continued and that (z'_n) has the claimed properties.

Now it is fairly easy to conclude the proof of the Josefson–Nissenzweig theorem. With (z'_n) as in the second claim we denote by $T : X \rightarrow \ell^\infty$ the operator $x \mapsto (z'_1(x), z'_2(x), \dots)$, and we define w'_n to be the functional $x \mapsto L(\lambda_n T x)$ (notation as in section 5) where L is a fixed Banach limit. Every w'_n has a norm not smaller than $\delta - \tau$ since $\lambda_n T x_n$ is by construction a sequence which – up to finitely many exceptions – is τ -close to the sequence $(\delta, \delta, \delta, \dots)$, and the Banach limit property of L implies that $L(\lambda_n T x_n)$ is – up to $\tau - L(\delta, \delta, \dots) = \delta$.

The w'_n also tend to zero w.r.t. the weak* topology since by 5.2 for every $x \in \ell^\infty$ with $\|x\| = 1$ the numbers $r_{\varepsilon_1 \dots \varepsilon_\ell} := L(\mu_{\varepsilon_1 \dots \varepsilon_\ell} x)$ satisfy the hypothesis of 5.1 and since the η_ℓ of 5.1 are just the $L(\lambda_\ell \cdot x)$ in this case.

Note. In fact we have shown more than required: When case 2.2 leads to a Josefson–Nissenzweig sequence then it is not only weak* null but in fact a weak*- ℓ^2 -sequence (i.e. $\sum |w'_n(x)|^2 < \infty$ for every $x \in X$).

REFERENCES

- [1] Behrends, E.: On Rosenthal's ℓ^1 -theorem
to appear (Archiv d. Math., 1994).
- [2] Diestel, J.: Sequences and Series in Banach Spaces
Springer-Verlag, Berlin-Heidelberg-New York, 1984.
- [3] Hagler, J. and Johnson, W.B.: On Banach spaces whose dual balls are
not weak* sequentially compact
Israel J. Math. 28, 1977, 325-330.
- [4] Josefson, B.: Weak sequential convergence in the dual of a Banach
space does not imply norm convergence
Ark. Mat. 13, 1975, 79-89.
- [5] Nissenzweig, A.: w^* sequential convergence
Israel J. Math. 22, 1975, 266-277.
- [6] Rosenthal, H.P.: A characterization of Banach spaces containing ℓ^1
Proc. Nat. Acad. Sci. (USA) 71, 1974, 2411-2413.

I. Mathematisches Institut
 FU Berlin
 Arnimallee 2-6
 D-14195 Berlin
 e-mail: behrends@math.fu-berlin.de